arXiv:1112.3153v2 [cs.DM] 22 Dec 2011

A new proof to the complexity of the dual basis of a type I optimal normal basis

Baofeng Wu*, Kai Zhou, Zhuojun Liu

Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China

---

Abstract

The complexity of the dual basis of a type I optimal normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ was determined to be $3n-3$ or $3n-2$ according as q is even or odd, respectively, by Z.-X. Wan and K. Zhou in 2007. We give a new proof to this result by clearly deriving the dual of a type I optimal basis with the aid of a lemma on the dual of a polynomial basis.

Keywords: Finite fields; Normal basis; Complexity; Type I optimal normal basis; Dual basis; Polynomial basis.

---

1. Introduction

Let $\mathbb{F}_{q^n}$ be an extension of the finite field $\mathbb{F}_q$. A basis of the form $\{\alpha, \alpha^q, \ldots, \alpha^{q^{n-1}}\}$ is called a normal basis, and α is called a normal basis generator in this case. Normal bases have great advantages against the common polynomial bases in arithmetic of finite fields. A simple property about a normal basis is it has normal dual basis. By dual basis $\{\beta_1^*, \ldots, \beta_n^*\}$ of a given basis $\{\beta_1, \ldots, \beta_n\}$, we mean $\mathrm{Tr}_{\mathbb{F}_{q^n}/\mathbb{F}_q}(\beta_i\beta_j^*) = \delta_{ij}$, $1 \leq i, j \leq n$.

For a normal basis generator α, denote $\alpha_i = \alpha^{q^i}$, $i = 0, \ldots, n-1$ and $N = \{\alpha_0, \ldots, \alpha_{n-1}\}$. Let

$$\alpha \cdot \alpha_i = \sum_{j=0}^{n-1} t_{ij}\alpha_j, \quad t_{ij} \in \mathbb{F}_q,\ 0 \leq i \leq n-1.$$

---

*Corresponding author

Email address: `wubaofeng@amss.ac.cn` (Baofeng Wu)

The number of nonzero elements in the $n \times n$ matrix $T = (t_{ij})$ is called the complexity of the normal basis N. We denote it by C_N. It can be shown that $C_N \geq 2n-1$, and N is called optimal when the lower bound is attained.

A type I optimal normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is the normal basis formed by the n nonunit $(n+1)$-th roots of unity, where $n+1$ is a prime and q is primitive modulo $n+1$ [2]. In [3] the dual of a type I optimal normal basis was studied, and its complexity was determined to be $3n-3$ or $3n-2$ according as q is even or odd, respectively.

Two bases $\{\beta_1, \ldots, \beta_n\}$ and $\{\beta'_1, \ldots, \beta'_n\}$ of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ are called equivalent, or weakly equivalent, if $\beta_i = c\beta'_i$, $i = 1, \ldots, n$ for some $c \in \mathbb{F}_q$, or $\mathbb{F}_{q^n}$, respectively. It is easy to see equivalent normal bases share the same complexity.

In this paper, we clearly derive the dual M of a type I optimal normal basis N through a new approach and thus rediscover the complexity of it. The main auxiliary lemma we use will be presented in section 2, and in section 3, we will propose our method to get M and its complexity, which is totally different from that in [3] and seems more simple and easier to understand.

2. Auxiliary lemma on the dual of a polynomial basis

The following lemma, proposed as an exercise in [1, Chapter 2], is the main result we will need in computing the dual of a type I optimal normal basis. We will include a short proof for it here for completeness.

Lemma 1. *Let $\mathbb{F}_{q^n} = \mathbb{F}_q(\alpha)$ be an extension of the finite field $\mathbb{F}_q$, and $f(x) \in \mathbb{F}_q[x]$ be the minimal polynomial of α. Assume*

$$\frac{f(x)}{x-\alpha} = \beta_0 + \beta_1 x + \cdots + \beta_{n-1}x^{n-1} \in \mathbb{F}_{q^n}[x].$$

Then the dual basis of the polynomial basis $\{1, \alpha, \ldots, \alpha^{n-1}\}$ is $\frac{1}{f'(\alpha)}\{\beta_0, \beta_1, \ldots, \beta_{n-1}\} := \{\frac{\beta_0}{f'(\alpha)}, \frac{\beta_1}{f'(\alpha)}, \ldots, \frac{\beta_{n-1}}{f'(\alpha)}\}$, where f' is the formal derivative of f.

Proof. Let σ_i be the i-th Frobenius automorphism of $\mathbb{F}_{q^n}/\mathbb{F}_q$ and $\alpha_i = \sigma_i(\alpha) = \alpha^{q^i}$, $0 \leq i \leq n-1$. Then $f(x) = \prod_{i=0}^{n-1}(x-\alpha_i)$. For $l = 0, 1, \ldots, n-1$, we construct the auxiliary polynomial

$$F_l(x) = \sum_{i=0}^{n-1} \frac{f(x)}{x-\alpha_i} \frac{\alpha_i^l}{f'(\alpha_i)} - x^l.$$

It's easy to check $F_l(\alpha_i) = 0$ for $i = 0, 1, \ldots, n-1$. Thus $F_l(x) \equiv 0$ since $\deg F_l(x) \leq n-1$. On the other hand,

$$\begin{aligned}
F_l(x) &= \sum_{i=0}^{n-1} \sigma_i\Big(\frac{f(x)}{x-\alpha}\frac{\alpha^l}{f'(\alpha)}\Big) \\
&= \sum_{i=0}^{n-1} \sigma_i\Big(\sum_{j=0}^{n-1} \beta_j x^j \frac{\alpha^l}{f'(\alpha)}\Big) \\
&= \sum_{j=0}^{n-1} \left(\sum_{i=0}^{n-1} \sigma_i\Big(\beta_j \frac{\alpha^l}{f'(\alpha)}\Big)\right) x^j \\
&= \sum_{j=0}^{n-1} \mathrm{Tr}_{\mathbb{F}_{q^n}/\mathbb{F}_q}\Big(\frac{\beta_j}{f'(\alpha)}\alpha^l\Big) x^j.
\end{aligned}$$

So we get $\mathrm{Tr}_{\mathbb{F}_{q^n}/\mathbb{F}_q}\big(\frac{\beta_j}{f'(\alpha)}\alpha^l\big) = \delta_{jl}$ for any $0 \leq j, l \leq n-1$, which is what we want to prove. $\square$

Remark 1. In fact, the lemma and the proof hold for any separable algebraic field extension $K(\alpha)/K$.

3. Dual basis of a type I optimal normal basis and its complexity

Recall that a type I optimal normal basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is of the form $N = \{\alpha, \alpha^2, \ldots, \alpha^n\} = \alpha\{1, \alpha, \ldots, \alpha^{n-1}\}$ where α is a root of the primitive polynomial $f(x) = \sum_{i=0}^n x^i$ [2], i.e. it is weakly equivalent to the polynomial basis $\{1, \alpha, \ldots, \alpha^{n-1}\}$. Thus if we can compute the dual of the polynomial basis, say $\frac{1}{f'(\alpha)}\{\beta_0, \beta_1, \ldots, \beta_{n-1}\}$ by Lemma 1, then we can get the dual of N of the form $M = \frac{1}{\alpha f'(\alpha)}\{\beta_0, \beta_1, \ldots, \beta_{n-1}\}$. The remaining task is just to compute $\alpha f'(\alpha)$ and $\beta_0, \ldots, \beta_{n-1}$.

Lemma 2.

$$\alpha f'(\alpha) = \frac{n+1}{\alpha - 1}.$$

Proof. Since $f(x) = \prod_{i=1}^n (x - \alpha^i)$, we know that

$$f'(\alpha) = \prod_{i=2}^{n} (\alpha - \alpha^i) = \alpha^{n-1} \prod_{i=1}^{n-1} (1 - \alpha^i) = \alpha^{n-1} \frac{f(1)}{1-\alpha^n} = \frac{\alpha^{n-1}}{1-\alpha^n}(n+1).$$

Then

$$\alpha f'(\alpha) = \frac{\alpha^n}{1-\alpha^n}(n+1) = \frac{\alpha^{n+1}}{\alpha-\alpha^{n+1}}(n+1) = \frac{n+1}{\alpha-1}.$$

$\square$

Lemma 3.

$$\beta_i = \frac{\alpha^{n-i}-1}{\alpha-1}, \quad i = 0, 1, \ldots, n-1.$$

Proof. Since $\frac{f(x)}{x-\alpha} = \prod_{i=2}^{n}(x-\alpha^i) = \sum_{i=0}^{n-1}\beta_i x^i$, we get $\beta_{n-1} = 1$ and $\beta_{n-i} = (-1)^{i-1}\sigma_{i-1}(\alpha^2, \alpha^3, \ldots, \alpha^n)$ for $i = 2, \ldots, n$, where σ_k stands for the k-th elementary symmetric polynomial.

As $f(x) = \prod_{i=1}^{n}(x-\alpha^i) = \sum_{i=0}^{n} x^i$, we know that

$$\begin{aligned} 1 &= (-1)^i\sigma_i(\alpha, \alpha^2, \alpha^3, \ldots, \alpha^n) \\ &= (-1)^i[\sigma_i(\alpha^2, \alpha^3, \ldots, \alpha^n) + \alpha\sigma_{i-1}(\alpha^2, \alpha^3, \ldots, \alpha^n)] \\ &= (-1)^i\sigma_i(\alpha^2, \alpha^3, \ldots, \alpha^n) - \alpha(-1)^{i-1}\sigma_{i-1}(\alpha^2, \alpha^3, \ldots, \alpha^n) \\ &= \beta_{n-i-1} - \alpha\beta_{n-i}. \end{aligned}$$

Thus

$$\begin{aligned} \beta_{n-i-1} &= 1 + \alpha\beta_{n-i} \\ &= 1 + \alpha(1 + \alpha\beta_{n-i+1}) \\ &= 1 + \alpha + \alpha^2\beta_{n-i+1} \\ &= \cdots \\ &= 1 + \alpha + \alpha^2 + \cdots + \alpha^i\beta_{n-1} \\ &= 1 + \alpha + \cdots + \alpha^i \\ &= \frac{\alpha^{i+1}-1}{\alpha-1} \end{aligned}$$

for $i = 1, \ldots, n-1$. So $\beta_i = \frac{\alpha^{n-i}-1}{\alpha-1}$ for $i = 0, 1, \ldots, n-1$. $\square$

From Lemma 2 and Lemma 3, we can clearly get that the dual of N is

$$M = \frac{1}{\alpha f'(\alpha)}\{\beta_0, \beta_1, \ldots, \beta_{n-1}\} = \frac{1}{n+1}\{\alpha^n - 1, \alpha^{n-1} - 1, \ldots, \alpha - 1\}.$$

Theorem 4. *The complexity of the dual of a type I optimal normal basis is* $3n-3$ *when* q *is even, and* $3n-2$ *when* q *is odd.*

Proof. We use the notations above. We only need to compute the complexity of $(n+1)M = \{\alpha^n - 1, \alpha^{n-1} - 1, \ldots, \alpha - 1\}$ as it is equivalent to M.

Since

$$\begin{aligned}(\alpha^n - 1)(\alpha^i - 1) &= \alpha^{n+i} - \alpha^n - \alpha^i + 1 \\ &= (\alpha^{i-1} - 1) - (\alpha^n - 1) - (\alpha^i - 1) \\ &= \begin{cases} -(\alpha^n - 1) - (\alpha - 1) & \text{if } i = 1 \\ (\alpha^{i-1} - 1) - (\alpha^n - 1) - (\alpha^i - 1) & \text{if } 2 \le i \le n-1 \\ (\alpha^{n-1} - 1) - 2(\alpha^n - 1) & \text{if } i = n, \end{cases}\end{aligned}$$

we can get the complexity of $(n+1)M$ is $2 + 3(n-2) + 1 = 3n - 3$ when q is even, and is $2 + 3(n-2) + 2 = 3n - 2$ when q is odd. □

Remark 2. Note that the main observation we make in deriving the dual of a type I optimal normal basis generated by α is the weak equivalence between it and the polynomial basis generated by α, so it is a natural question that when will a normal basis generator β generate a normal basis weakly equivalent to the polynomial basis generated by it in a general finite field $\mathbb{F}_{q^n}$. We remark that this happens if and only if β is a type I optimal normal basis generator. The reason is simple: if there exists some $\gamma \in \mathbb{F}_{q^n}$ such that $N = \{\beta, \beta^q, \ldots, \beta^{q^{n-1}}\} = \gamma\{1, \beta, \ldots, \beta^{n-1}\}$, we know that $\beta = \gamma\beta^k$ for some $k \in \mathbb{Z}$, $0 \le k \le n-1$. Thus $\gamma = \beta^{1-k}$. If $k \ne 0$, $\gamma\beta^{k-1} = \beta^0 = 1$, which is impossible since it is an element of N. So $\gamma = \beta$. A normal basis of the form $\{\beta, \beta^2, \ldots, \beta^n\}$ must be a type I optimal normal basis, as its complexity is no more than $2n-1$.

References

[1] R. Lidl, H. Niederreiter, Finite Fields, second ed., Encyclopedia Math. Appl., vol. 20, Cambridge University Press, Cambridge, 1997.

[2] A.J. Menezes, I.F. Blake, X.H. Gao, R.C. Mullin, S.A. Vanstone, T. Yaghoobian, Applications of Finite Fields, Kluwer Academic, Boston, MA, 1993.

[3] Z.-X. Wan, K. Zhou, On the complexity of the dual basis of a type I optimal normal basis, Finite Fields Appl. 13(2007) 411-417.